






Basal Eudicots

... transition from basal angiosperm
to advanced eudicot ...

Basal Eudicots




- already looked at basal angiosperms except monocots
- Eudicots are the majority of angiosperms and defined by 3 pored pollen - often called tricolpates




Basal Eudicots

- tricolpate pollen: only morphological feature defining eudicots consistently




Basal Eudicots




- tricolpate pollen: a derived or advanced character state that has evolved essentially once

- selective advantage for pollen germination




Basal Eudicots



- basal eudicots are a grade at the base of eudicots - paraphyletic
- morphologically are transitional between basal angiosperms and the core eudicots



Basal Eudicots



- basal eudicots are a grade at the base of eudicots - paraphyletic
- morphologically are transitional between basal angiosperms and the core eudicots
- examine two orders only:
Ranunculales - 7 families
Proteales - 3 families

Basal Eudicots

*Ranunculaceae (Ranunculales) - buttercup family

- largest family of the basal eudicots
- 60 genera, 2500 species
- distribution centered in temperate and cold regions of the northern and southern hemispheres



Basal Eudicots

*Ranunculaceae (Ranunculales) - buttercup family

- perennial herbs, sometimes woody or herbaceous climbers or low shrubs



Basal Eudicots

*Ranunculaceae (Ranunculales) - buttercup family

- perennial herbs, sometimes woody or herbaceous climbers or low shrubs
- leaves, alternate sheathing, usually basal and cauline, often divided or compound, or palmately lobed; no stipules.



Basal Eudicots

*Ranunculaceae (Ranunculales) - CA 3+ CO (0)5+ A ∞ G (1)3+ buttercup family

- floral diversity enormous except ∞ stamens and ∞ separate carpels
- insect (bird) pollination! nectar or pollen reward, radial or bilateral symmetry



Basal Eudicots

*Ranunculaceae (Ranunculales) - diverse fruit types buttercup family

Follicles = ∞ seeded dehiscent fruit



columbine

Berries = ∞ seeded fleshy fruit



baneberry

Achenes = 1 seeded indehiscent, dry fruit



buttercup

Basal Eudicots

*Ranunculaceae (Ranunculales) - buttercup family



Aconitum columbianum - monk's hood




Actaea rubra - red baneberry

Basal Eudicots

*Ranunculaceae (Ranunculales) -
buttercup family

only sepals!



Anemone patens - pasque flower

Anemonella thalictroides - rue anemone

Basal Eudicots

*Ranunculaceae (Ranunculales) -
buttercup family

Aquilegia canadensis - american columbine



Basal Eudicots

*Ranunculaceae (Ranunculales) -
buttercup family

Caltha palustris - marsh marigold



Basal Eudicots

*Ranunculaceae (Ranunculales) -
buttercup family

Sepals + petals, achenes

Ranunculus - buttercup, crowfoot



Basal Eudicots

*Ranunculaceae (Ranunculales) -
buttercup family

Dioecious - with separate male and female plants,
wind pollinated!



Thalictrum dioicum - early meadow-rue

Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

- 13 genera, 660 species; widespread in temperate regions of Northern hemisphere - ArctoTertiary relict distribution
- small shrubs (*Berberis*) or herbs (rest of family).



Podophyllum - mayapple



Berberis - barberry

Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

- all genera attacked by same family of rusts

Podophyllum – mayapple
with *Puccinia podophylli*



Podophyllum - mayapple




Berberis - barberry

Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

CA 6 CO 6/9 A 6/9 G 1

- 3 merous flowers, 1 superior carpel, marginal placentation, berry



Podophyllum peltatum - mayapple



Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

Podophyllum hexandrum
eastern Asia



Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

Berberis vulgaris
European barberry



Berberis thunbergii
Japanese barberry



Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

Caulophyllum thalictroides
Blue cohosh



Basal Eudicots

*Berberidaceae (Ranunculales) -
barberry family

4-merous!

Jeffersonia diphylla - twinleaf

J. dubia
Eastern Asia



Basal Eudicots

Menispermaceae (Ranunculales) -
moonseed family

- mainly tropical family, arrow poisons, usually viney
- dioecious; separate carpels each produce one seeded drupes
= fleshy with one bony seed



Chondrodendron curare



Menispermum canadense moonseed



Basal Eudicots

*Papaveraceae (Ranunculales) -
poppy family

CA 2 CO 4/8 A ∞ G (2- ∞)

- 2 deciduous sepals, petals crumpled in bud



celadine



poppy

Basal Eudicots

*Papaveraceae (Ranunculales) -
poppy family

- 25 genera and about 200 species mainly of north temperates
- herbaceous, alternate compound leaves with colored latex, narcotics



Papaveraceae



celadine



poppy

Basal Eudicots

*Papaveraceae (Ranunculales) -
poppy family

Papaver somniferum
opium poppy

CA 2 CO 4/8 A ∞ G (2- ∞)

- pistil syncarpic forming capsule = dehiscent syncarpic fruit



celadine



Basal Eudicots

*Papaveraceae (Ranunculales) -
poppy family



Sanguinaria canadensis - bloodroot

Basal Eudicots

*Papaveraceae (Ranunculales) -
poppy family



Eschscholzia - California poppy

Stylophorum - celadine poppy

Basal Eudicots

*Papaveraceae (Ranunculales) -
poppy family

- family Fumariaceae now in Papaveraceae = zygomorphic poppies with 2 carpels



Dicentra cucullaria - Dutchman's breeches

2 carpellate capsules

Basal Eudicots


*Papaveraceae (Ranunculales) -
poppy family

- family Fumariaceae now in Papaveraceae = zygomorphic poppies with 2 carpels



Corydalis

Basal Eudicots




- Proteales - unusual order of 3 families
 - Nelumbonaceae - lotus lily family
 - Platanaceae - sycamore family
 - Proteaceae - macadamia family
- all 3 share unusual fruiting structures - 1 carpellate fruits in “heads”



Basal Eudicots

Nelumbonaceae (Proteales) - lotus lily family




one genus and two species - one North American and the second East Asian (sacred oriental lotus lily)

Nelumbo lutea - lotus lily
Nymphaea odorata - water lily

once considered related to water lilies of Nymphaeaceae - now known to be convergence

Basal Eudicots

Nelumbonaceae (Proteales) - lotus lily family



one genus and two species - one North American and the second East Asian (sacred oriental lotus lily)




Nelumbo lutea - lotus lily

Habit is an aquatic, floating or emergent leaved, perennial herb. Leaves are simple, peltate. Inflorescence is a solitary flower.

Basal Eudicots

Nelumbonaceae (Proteales) - lotus lily family

P ∞ A ∞ G [∞]




Nelumbo lutea - lotus lily

Basal Eudicots

Nelumbonaceae (Proteales) -
lotus lily family

- separate, superior one-seeded carpels but held in the cavities of the obconical receptacle



Nelumbo lutea - lotus lily

Basal Eudicots

Nelumbonaceae (Proteales) -
lotus lily family

- “fruit” is a woody receptacle cone with a nut-like seed laying loose in each cavity



Nelumbo lutea - lotus lily

Basal Eudicots

Platanaceae (Proteales) - sycamores

- 1 genus with 10 species of northern hemisphere trees, distinctive bark

Platanus occidentalis - sycamore



Basal Eudicots

Platanaceae (Proteales) - sycamores


- monoecious, female flowers apocarpic in spherical heads - wind pollination
- fruit = head of achenes (one seeded)
- achene with coma - ant dispersed



Basal Eudicots

Platanaceae (Proteales) - sycamores

- European plane tree - *Platanus x hybrida* - important urban shade tree



Basal Eudicots

Proteaceae (Proteales) - proteas

- large southern hemisphere family of 75 genera, 1400 species - ecologically and morphologically diverse

- inflorescences of 4 merous flowers with single superior carpel



Basal Eudicots

Proteaceae (Proteales) - proteas

- large southern hemisphere family of 75 genera, 1400 species - ecologically and morphologically diverse

- infructescence - often woody "cone" with single seeded fruits; serotinous

